

2. (Currently Amended) ~~A The process according to claim 1, wherein dried playhcodi platycodi radix (broad bellflower), ginseng radix (Panax ginseng) and peach kernel (persicae semen) are crushed, are mixed with cut eum-yang-kwak (dried leaves of epimedii herbal tea (*Epimedium koreanum*)) are cut into the same size as that of the cut tobacco leaves and mixture of playcodi radix (broad bellflower), ginseng radix (*Panax ginseng*) and peach kernel (persicae semen), eum-yang-kwak (dried leaves of epimedii herbal tea (*Epimedium koreanum*)) in the same proportion~~ equal proportions before being applied to the tobacco leaves.

3. (Currently Amended) ~~A The process according to claim 1, wherein the mixture proportion of dried playhcodi platycodi radix (broad bellflower), ginseng radix (Panax ginseng) and peach kernel (persicae semen) into to tobacco leaves is at the a ratio of 3:3:3:991.~~

4. (Currently Amended) ~~A The process according to claim 1, wherein playcodi the platycodi radix (broard broad bellflower) harvested are 2 ~ 3 years old.~~

5. (Currently Amended) ~~A The process according to claim 1, wherein the ginseng radix (Panax ginseng) harvested is 1 year old.~~

6. (Currently Amended) ~~A The process according to claim 1, wherein the peach kernel (persicae semen) is harvested in June ~ August.~~

7. (Currently Amended) ~~★~~ The process according to claim 1, wherein the mixture of dried ~~playcodi~~ ~~platycodi~~ radix (broad bellflower), ginseng radix (Panax ginseng) and peach kernel (persicae semen) into tobacco leaves are at ~~the~~ a ratio of 3:3:3 ~ 5:989 ~ 991.

8. (Currently Amended) ~~★~~ The process according to claim 1, wherein glycerin is sprayed slightly on the crushed dried ~~playcodi~~ ~~platycodi~~ radix (broad bellflower), ginseng radix (Panax ginseng) and peach kernel (persicae semen) and ~~eum-yang-kwak (dried leaves of epimedium~~ herbal tea (*Epimedium koreanum*)).

9. (Currently Amended) ~~★~~ The process according to claim 1, wherein dried ~~playcodi~~ ~~platycodi~~ radix (broad bellflower), ginseng radix (Panax ginseng) and peach kernel (persicae semen) are crushed to a size, which ~~would not enable them to size does not~~ infiltrate through a cigarette filter.

10. (Currently Amended) ~~★~~ The process according to claim 1, wherein dried ~~playcodi~~ ~~platycodi~~ radix (broad bellflower), ginseng radix (Panax ginseng) and peach kernel (persicae semen) are crushed to a size of ~~100 ~ 130 meshes between 100 to 130 mesh.~~

11. (Currently Amended) ~~★~~ The process according to claim 1, wherein ~~playcodi~~ ~~platycodi~~ radix (broad bellflower), ginseng radix (Panax ginseng) and peach kernel (persicae semen) and ~~eum-yang-kwak (dried leaves of epimedium~~ herbal tea (*Epimedium koreanum*)) are dried in the shade.

12. (Currently Amended) ~~A~~ The process according to claim 2, wherein ~~platycodi~~ ~~the amount of platycodi~~ radix (broad bellflower), ginseng radix (*Panax ginseng*) and peach kernel (*Persicae semen*) ~~are added is $\pm 0.1\%$~~ , and tobacco leaves and eum-yang-kwak (dried leaves of *Epimedium koreanum*) which were cut into a specified size are mixed at ~~the~~ a ratio of 6:4.

13. (Currently Amended) ~~A~~ The process according to claim 11, wherein ~~platycodi~~ platycodi radix (broad bellflower), ginseng radix (*Panax ginseng*) and peach kernel (*Persicae semen*) and eum-yang-kwak (dried leaves of *Epimedium koreanum*) are dried to ~~the have a~~ water content of as follows: platycodi radix (broad bellflower) is 6 ~ 10% , ginseng radix (*Panax ginseng*) 10-14%, peach kernel (*Persicae semen*) 6 ~ 10% and eum-yang-kwak (dried leaves of *Epimedium koreanum*) 3 ~ 7%.

14. (Previously Presented) ~~A~~ The process according to claim 2, wherein dried ~~platycodi~~ platycodi radix (broad bellflower), ginseng radix (*Panax ginseng*) and peach kernel (*Persicae semen*) are crushed to a size, which ~~would not enable them to~~ size does not infiltrate through a cigarette filter.

15. (Previously Presented) ~~A~~ The process according to claim 2, wherein dried ~~platycodi~~ platycodi radix (broad bellflower), ginseng radix (*Panax ginseng*) and peach kernel (*Persicae semen*) are crushed to a size of ~~100-130 meshes~~ between 100 to 130 mesh.

16. (Previously Presented) ~~A~~ The process according to claim 2, wherein platycodi platycodi radix (broad bellflower), ginseng radix (*Panax ginseng*) and peach kernel (*Persicae semen*) and eum-yang-kwak (dried leaves of *Epimedium koreanum*) are dried in the shade.

17. (Currently Amended) ~~A~~ The process according to claim 16, wherein platycodi platycodi radix (broad bellflower), ginseng radix (*Panax ginseng*) and peach kernel (*Persicae semen*) and eum-yang-kwak (dried leaves of *Epimedium koreanum*) are dried to ~~the~~ have a water content of as follows: platycodi radix (broad bellflower) is 6-10%, ginseng radix (*Panax ginseng*) 10-14%, peach kernel (*Persicae semen*) 6-10% and eum-yang-kwak (dried leaves of *Epimedium koreanum*) 3-7%.